

Lung function corner

Putting lung function reference equations into context

Case history

A 69-year-old man with idiopathic pulmonary fibrosis (IPF) is undertaking routine lung function testing prior to clinical review and assessment for consideration of antifibrotics. As he lives in Australia, the pulmonary function eligibility criteria for antifibrotics are forced vital capacity (FVC) $\geq 50\%$ of predicted and transfer factor of the lung for carbon monoxide (T_{LCO}) $\geq 30\%$ of predicted in the absence of obstruction [1].

Questions

Does the patient meet the lung function criteria for antifibrotics based on the lung function results in table 1?

Are the reference equations used appropriate for this patient?

Answers

Table 1 shows the lung function results of spirometry, static lung volumes (SLVs) and T_{LCO} . Results show a restrictive ventilatory defect as evidenced by a reduced total lung capacity (below the lower limit of normal (LLN) or fifth centile) in the presence of no evidence of obstruction on spirometry (forced expiratory volume in 1 s (FEV_1)/FVC $>$ LLN or fifth centile). This is in keeping

with the patient's known IPF. T_{LCO} unadjusted for haemoglobin, alveolar volume and the transfer coefficient are all reduced ($<$ LLN or fifth centile), revealing gas exchange impairment suggestive of parenchymal or pulmonary vascular disease. When corrected for haemoglobin, gas exchange ($T_{\text{LCO}_{\text{Hbcorr}}}$) remains impaired.

The patient's FVC is 48% of the predicted value and T_{LCO} uncorrected for haemoglobin is 29% of the predicted value. Based on these results, the patient does not meet the eligibility criteria for antifibrotics. However, $T_{\text{LCO}_{\text{Hbcorr}}}$ is 30% of the predicted value and if used, would meet the T_{LCO} criterion.

The patient's ethnicity is documented on the report as "other/mixed". This means he does not identify as one of the four ethnic groups assigned in the Global Lung Function Initiative (GLI) spirometry reference sets. The reference sets used (documented at the bottom of the report) (table 1) are stated as GLI spirometry Caucasian [2], GLI SLVs [3] and GLI T_{LCO} [4]. As the patient identifies as a race other than Caucasian or is of mixed race, the selected spirometry reference set is not appropriate for this patient. Furthermore, the GLI SLV and GLI T_{LCO} reference sets are based on Caucasian data and may also be inappropriate for this patient.

Discussion

The selection of appropriate reference sets is critical to interpretation of results. Incorrect selection of

Cite as: Borg BM, Thompson BR. Putting lung function reference equations into context. *Breathe* 2021; 17: 210099.

 @ERSpublications

Knowing the limitations of reference equations is essential to minimising errors in diagnosis and clinical management. Choice of reference sets may impact access to treatment options where lung function based eligibility criteria exist. <https://bit.ly/2WdOFDj>



CrossMark



© ERS 2021

Table 1 Lung function test results

Gender	Male	Age, years	69.8	Race	Other/mixed
Height, cm	171.8	Weight, kg	96	BMI, kg·m⁻²	33
Clinical notes	Idiopathic pulmonary fibrosis				
	Actual	LLN	ULN	Z-score	% pred
Pre-bronchodilator spirometry					
FVC, L	1.89	2.90		-3.31	48
FEV ₁ , L	1.53	2.15		-2.77	52
FEV ₁ /FVC	0.81	0.63		0.62	
SLVs					
TLC, L	2.48	5.22	7.99	-5.06	37
VC, L	1.87	3.38		-4.69	44
FRC, L	1.39	2.43	4.83	-3.809	
RV, L	0.61		3.46	-3.44	
RV/TLC, %	25		47	-1.59	
CO gas transfer					
IVC, L	1.77				
V _A , L	2.50	4.76		-5.41	
T _{LCO} , mmol·min ⁻¹ ·kPa ⁻¹	2.31	5.95	10.6	-5.63	29
T _{LCOHbcorr} , mmol·min ⁻¹ ·kPa ⁻¹	2.45			-5.42	30
K _{CO} , mmol·min ⁻¹ ·kPa ⁻¹ ·L ⁻¹	0.92	1.02	1.76	-2.13	
K _{COHbcorr} , mmol·min ⁻¹ ·kPa ⁻¹ ·L ⁻¹	0.98			-1.86	
Hb, g·dL ⁻¹	12.6	12.2	17.0		
Technical comment	Grade: spirometry: FEV ₁ – A, FVC – A; SLV: good; T _{LCO} : AA Ref: GLI spirometry Caucasian, GLI T _{LCO} , GLI SLV				

Results show a restrictive ventilatory defect with gas exchange impairment suggestive of lung parenchymal or pulmonary vascular disease. Reference sets used (Ref) may not be representative of the patient's ethnicity and should be used with caution. BMI: body mass index; LLN: lower limit of normal; ULN: upper limit of normal; FVC: forced vital capacity; FEV₁: forced expiratory volume in 1 s; SLV: static lung volume; TLC: total lung capacity; VC: vital capacity; FRC: functional residual capacity; RV: residual volume; IVC: inspiratory vital capacity; V_A: alveolar volume; T_{LCO}: transfer across the lung for carbon monoxide; Hbcorr: corrected for haemoglobin; K_{CO}: transfer coefficient of the lung for carbon monoxide; Hb: Haemoglobin; GLI: Global Lung Function Initiative [2–4].

reference equations may impact diagnosis and incorrectly inform clinical management options.

The multiethnic GLI spirometry reference set illustrates differences in lung function based on ethnicity, with Caucasian and North East Asian populations having higher FEV₁ and FVC predicted values than South East Asian and African American populations in the study [2]. Figure 1 illustrates the differences. Of note, in the examples shown in figure 1, the North East Asian LLN sits higher than the Caucasian LLN, despite the Caucasian predicted value being higher than the North East Asian predicted value. While this suggests less variation in the North East Asian population used to create the reference equations, it also suggests that using Caucasian reference equations in a North East Asian population may result in underestimation

of abnormal findings. Using Caucasian reference values for a person of South East Asian background may result in an abnormal finding simply because the LLN using the Caucasian reference equations is higher than the South East Asian LLN (table 2). Similarly, a person of Caucasian descent may appear to have lung function in the normal range if compared against the South East Asian LLN but be abnormal if compared to the Caucasian LLN.

In the case presented, the subject has been assigned “other/mixed” for ethnic ancestry as he does not identify with the four defined GLI ethnic groupings or is of mixed ancestry. The GLI “other/mixed” equations represent the combination of the entire data set and hence sit somewhere in the middle of the range (figure 1). Cautionary statements regarding reference equations must be




Figure 1 a) Predicted (pred) FVC and FVC LLN for a 172-cm-tall adult male versus age using the multiethnic GLI equations. b) Pred FVC and FVC LLN for a 163-cm-tall adult female. For both, the “other/mixed” pred FVC and FVC LLN represent the pooled data of the four ethnic groups and sit approximately in the middle of the range of plotted data. Note also that the North East (NE) Asian FVC LLN sits above the Caucasian FVC LLN despite the NE Asian predicted FVC sitting below the Caucasian pred FVC. SE: South East.

used when “other/mixed” group is selected, as it is unclear whether the reference values are truly representative of the subject’s ethnicity and they may be assigned normal or abnormal patterns based on a generalised normal range.

While the GLI spirometry reference set, which has been endorsed for use by many respiratory societies around the world, provides equations for multiple ethnic groups, similar multiethnic reference equations are not available for T_{LCO} or SLV parameters. Whether similar differences in gas exchange and SLV parameters across racial groups exist is still to be determined. However, it is probable that similar findings of racial differences for SLVs exist based on the common parameter of vital capacity (VC). In a normal, healthy cohort without any history of respiratory disease, we expect FVC and VC to be similar. Spirometry data submitted with lung volume data used to create the GLI SLV reference set overall showed small, insignificant deviations from the GLI predicted FVC (mean±SD z-score 0.31 ± 1.00) [3]. Furthermore, equations for VC [3] and FVC [2] were similar [3]. The differences between VC and FVC amplify, however, when using spirometry reference equations other than Caucasian to calculate FVC. Figure 2 shows that the difference between predicted FVC and predicted VC is larger when predicted FVC is calculated using reference equations appropriate to ancestry in a male cohort (n=365) referred to an Australian lung function laboratory for clinical testing (unpublished data). By assigning the “other/mixed” spirometry reference set to the case presented, the predicted FVC and LLN become 3.60 L and 2.72 L respectively, while the predicted VC and LLN are 4.25 L and 3.38 L respectively: a 0.65 L and 0.66 L difference respectively. It is probable that the difference seen between predicted VC and FVC in this case is likely to be reflected in other subdivisions of SLVs too. SLV reference data for ancestry other than Caucasian are sparse and it is unclear how to account for possible differences when using Caucasian reference

equations to interpret results of patients with ancestry other than Caucasian. It is incumbent upon reporters of lung function to understand the limitations of the reference sets being used; the use of cautionary statements in reports is strongly recommended where the ancestry of the patient differs from the reference sets used (e.g. reference values used may not be representative of the patient’s ethnicity and should be used with caution).

Table 2 Comparison of spirometry results of a 21-year-old, 163-cm-tall female of South East Asian ancestry using GLI South East Asian and Caucasian reference values [2]

	Measured	GLI reference set			
		South East Asian		Caucasian	
		LLN	Z-score	LLN	Z-score
FEV₁, L	2.04	1.81	-1.02	2.09	-1.78
FVC, L	2.56	2.24	-0.89	2.65	-1.83
FEV₁/FVC	0.80	0.72	-0.42	0.68	-0.02




Figure 2 Male predicted FVC [2] minus predicted VC [3] plotted against age. “Non-Caucasian” includes North East Asian, South East Asian, African American or other/mixed ancestry. The difference in predicted FVC and VC is substantially larger in those of non-Caucasian ancestry.

Table 3 Comparison of FVC and T_{LCO} % predicted (pred) across reference sets for a 69.8-year-old, 171.8-cm-tall male of “other/mixed” ancestry as per the GLI with measured FVC 1.89 L and T_{LCO} 2.45 $\text{mmol}\cdot\text{min}^{-1}\cdot\text{kPa}^{-1}$

	FVC pred, L	FVC, % pred
GLI other/mixed [2]	3.60	53
GLI Caucasian [2]	3.91	48
ECSC [6]	3.80	50
KNUDSON <i>et al.</i> [9]	3.73	51
CRAPO <i>et al.</i> [10]	4.23	45
HANKINSON <i>et al.</i> [7], Caucasian	4.11	46
	T_{LCO} pred, $\text{mmol}\cdot\text{min}^{-1}\cdot\text{kPa}^{-1}$	T_{LCO} , % pred
GLI 2017 [4]	8.09	30
MILLER <i>et al.</i> [11]	7.84	31
ROCA <i>et al.</i> [12]	8.56	29
THOMPSON <i>et al.</i> [13]	7.76	32

ECSC: European Community for Steel and Coal.

Self-evaluation questions

- Which GLI reference equations provide multiethnic capability?
 - SLV
 - Spirometry
 - T_{LCO}
 - All the above
- Which spirometry GLI reference equations apply when the ethnic origin is not known?
 - Caucasian
 - Caucasian \times 0.88
 - South East Asian
 - African American
 - Other/mixed
- The “other/mixed” GLI spirometry reference equations are derived from:
 - Submitted data of individuals who did not identify as Caucasian, North East Asian, South East Asian or African American
 - All the GLI spirometry data combined
 - Caucasian and African American datasets only
 - North East and South East Asian datasets only

study was FVC \leq 80% of predicted. The eligibility criteria did not state which reference values should be used. Approximately 34 out of 98 patients (total $n=528$) who were ineligible for treatment using the European Community for Steel and Coal reference set [6] to calculate FVC % predicted became eligible when using the Third National Health and Nutrition Examination Survey spirometry reference set [7], and 25 became eligible using the GLI 2012 reference set [2] instead. The second study from Australia, where antifibrotic lung function eligibility criteria is FVC \geq 50% predicted and T_{LCO} \geq 30% predicted, found variation in eligibility ranged from 74% to 86% dependent on the reference sets chosen [8]. Table 3 shows the variation in FVC % predicted and T_{LCO} % predicted in a selection of commonly used reference sets using the presented case’s gender, age and height. Using the GLI spirometry “other/mixed” and T_{LCO} reference equations, the patient (just) meets the lung function eligibility criteria for government-subsidised treatment.

Conclusion

In the case presented, the patient has IPF and the medical team are assessing his eligibility to access antifibrotics. Quite apart from the differences in eligibility between jurisdictions around the world, the patient’s ability to access the medication may depend on the reference set chosen. Two recent papers have reviewed eligibility for access to antifibrotics based on reference set chosen. WARD *et al.* [5] looked at the proportion of English and Scottish patients with IPF who were eligible for antifibrotics dependent on reference sets chosen. Eligibility in England and Scotland at the time of the

It is essential for persons reporting lung function to know the reference sets being used and understand the limitations reference sets may have. Consideration of the impact of using reference sets that may not be representative of the patient’s ethnicity or race is necessary and cautionary statements should be used in lung function reports where this occurs. The choice of reference set may impact the eligibility for patients to access government-subsidised medications where lung function criteria are based on percent of predicted value.

Key points

- Knowing the limitations of lung function reference sets and understanding the impact of the limitations are essential to interpretation of lung function data.
- The multiethnic GLL spirometry reference equations demonstrate clear differences between groups. It is not clear how to use the Caucasian-based GLL T_{LCO} and SLV reference sets in people not identifying as Caucasian.
- Use cautionary statements to identify limitations of reference sets in reports.
- Choice of reference sets may impact patients seeking access to treatment with lung function related eligibility criteria.

Affiliations

Brigitte M. Borg^{1,2}, Bruce R. Thompson³

¹Respiratory Medicine, The Alfred, Melbourne, Australia. ²School of Public Health and Preventive Medicine, Monash University, Melbourne, Australia. ³School of Health Sciences, Swinburne University of Technology, Hawthorn, Australia.

Conflict of interest

B.M. Borg is co-author of a book *Interpreting Lung Function Tests: A Step-by-Step Guide* for which they receive royalties (Wiley-Blackwell Publishing). B.R. Thompson is co-author of a book *Interpreting Lung Function Tests: A Step-by-Step Guide* for which they receive royalties (Wiley-Blackwell Publishing).

References

1. Jo HE, Prasad JD, Troy LK, *et al.* Diagnosis and management of idiopathic pulmonary fibrosis: Thoracic Society of Australia and New Zealand and Lung Foundation Australia position statements summary. *Med J Aust* 2018; 208: 82–88.
2. Quanjer PH, Stanojevic S, Cole TJ, *et al.* Multi-ethnic reference values for spirometry for the 3–95-yr age range: the global lung function 2012 equations. *Eur Respir J* 2012; 40: 1324–1343.
3. Hall GL, Filipow N, Ruppel G, *et al.* Official ERS technical standard: Global Lung Function Initiative reference values for static lung volumes in individuals of European ancestry. *Eur Respir J* 2021; 57: 2000289.
4. Stanojevic S, Graham BL, Cooper BG, *et al.* Official ERS technical standards: Global Lung Function Initiative reference values for the carbon monoxide transfer factor for Caucasians. *Eur Respir J* 2017; 50: 1700010.
5. Ward K, Spurr L, Goldman NR, *et al.* Patient eligibility for anti-fibrotic therapy in idiopathic pulmonary fibrosis can be altered by use of different sets of reference values for calculation of FVC percent predicted. *Respir Med* 2016; 120: 131–133.
6. Quanjer PH, Tammeling GJ, Cotes JE, *et al.* Lung volumes and forced ventilatory flows. Report Working Party Standardization of Lung Function Tests, European Community for Steel and Coal. Official Statement of the European Respiratory Society. *Eur Respir J* 1993; 6: Suppl. 16, 5–40.
7. Hankinson JL, Odencrantz JR, Fedan KB. Spirometric reference values from a sample of the general U.S. population. *Am J Respir Crit Care Med* 1999; 159: 179–187.
8. Burgess A, Goon K, Brannan JD, *et al.* Eligibility for anti-fibrotic treatment in idiopathic pulmonary fibrosis depends on the predictive equation used for pulmonary function testing. *Respirology* 2019; 24: 988–995.
9. Knudson RJ, Burrows B, Lebowitz MD. The maximal expiratory flow-volume curve: its use in the detection of ventilatory abnormalities in a population study. *Am Rev Respir Dis* 1976; 114: 871–879.
10. Crapo RO, Morris AH, Gardner RM. Reference spirometric values using techniques and equipment that meet ATS recommendations. *Am Rev Respir Dis* 1981; 123: 659–664.
11. Miller A, Thornton JC, Warshaw R, *et al.* Single breath diffusing capacity in a representative sample of the population of Michigan, a large industrial state. Predicted values, lower limits of normal, and frequencies of abnormality by smoking history. *Am Rev Respir Dis* 1983; 127: 270–277.
12. Roca J, Rodriguez-Roisin R, Cobo E, *et al.* Single-breath carbon monoxide diffusing capacity prediction equations from a Mediterranean population. *Am Rev Respir Dis* 1990; 141: 1026–1032.
13. Thompson BR, Johns DP, Bailey M, *et al.* Prediction equations for single breath diffusing capacity (Tlco) in a middle aged caucasian population. *Thorax* 2008; 63: 889–893.

Suggested answers

1. b.
2. e.
3. b.